


What is claimed is:



1. A method of administering a compound of the formula













5 wherein X and Y are the same or different halo group; R is H or lower alkyl; and pharmaceutically acceptable salts thereof; together with a chemotherapeutic agent.




10 2. A method according to Claim 1 wherein the chemotherapeutic agent is selected from the group consisting of cisplatin; cyclophosphamide; 5-fluorouracil, doxorubicin and taxol.

15 3. A method according to Claim 1 wherein the compound is selected from the group consisting of:











5



4. A method according to Claim 3 wherein the chemotherapeutic agent
10 is selected from the group consisting of cisplatin,
cyclophosphamide, 5-fluorouracil, doxorubicin and taxol.

5. A method of treating or preventing a neoplasia disease comprising administering to a mammal in need of treatment for a neoplasia disease a therapeutically effective amount of a compound of the formula:



5



wherein X and Y are the same or different halo group; R is H or lower alkyl; or a pharmaceutically acceptable salt thereof; together with a chemotherapeutic agent.

10



6. A method according to Claim 5 wherein the chemotherapeutic agent is selected from the group consisting of cisplatin, cyclophosphamide, 5-fluorouracil, doxorubicin and taxol.

15




7. A method according to Claim 5 wherein the compound is selected from the group consisting of






20





5







10



5





5

8. A method according to Claim 7 wherein the chemotherapeutic agent
10 is selected from the group consisting of cisplatin,
cyclophosphamide, 5-fluorouracil, doxorubicin and taxol.
9. A pharmaceutical composition comprising a therapeutically effective
15 amount of a compound according to Claim 1, a chemotherapeutic
agent and a pharmaceutically acceptable carrier.

10. A method according to Claim 5 wherein the neoplasia disease is tumor metastasis.
11. A method according to Claim 5 wherein the neoplasia disease treated is solid tumor growth.
5
12. A method according to Claim 5 wherein the condition treated is angiogenesis.
- 10 13. A method according to Claim 5 wherein the neoplasia disease is humoral hypercalcemia of malignancy.